NOTES. 191

larva in extending the domain of a sea-dwelling animal, for, so far as I have been able to learn, no adults of this animal have been introduced into the coastal waters of either Sussex or Hampshire.

J. H. ORTON.

Note on the Land and Freshwater Shells of Texel and Terschelling. (*Read 13th November*, 1914.)—When paying a visit to these islands in August, 1913, the following species of land and freshwater Mollusca were collected.

Texel seemed fairly productive of shells, twenty species being found there. I was interested to find a Vertigo, none of which genus had

appeared in my exploration of Friesland some years ago.

Helicella cantiona, too, turned up unexpectedly, and could not, I think, have been imported, as I found it at Koog on the west side of the island, far from any harbour or place where boats land. Whether imported or not, it seems to have established itself in some numbers.

I also visited the intermediate island, Vlieland, but found no signs of

Mollusca there.

## Texel.

Helix nemoralis, L.
Pyranidula rotundata (Müll.).
Theba cantiana (Mont.), var. cantianiformis, Ancey.
Hygromia hispida (L.), var. hispidosa,
Mouss.
Cochlicopa lubrica (Müll.).
Vallonia excentrica, Sterki.
Vertigo antivertigo (Drap.).
Limnæa pereger (Müll.).

var. maritima, Jeff.
var. balthica.
var. acuminata, Jeff.
var. ovata, Drap.
Limnæa palustris (Müll.).
var. lacunosa (Zgl.).
Limnæa truncatula (Müll.).
Planorbis umbilicatus, Müll.
P. spirorbis (L.).

P. contortus (L.).
P. albus, Müll.
Succinea elegans, Risso, var. longiscata, Morelet.
Spherium corneum (L.).
Paludestrina ventrosa (Mont.).
Pisidium nitidum, Jenyns.
P. obtusale, Pfr.

## TERSCHELLING.

Limnæa pereger (Müll.).
L. palustris (Müll.).
Planorbis spirorbis (L.).
Sphærium corneum (L.).
Physa fontinalis (L.).
Succinea putris (L.).
Paludestrina stagnalis, Bast.
Pisidium casertanum, Poli.

P. subtruncatum, Malm.

F. H. SIKES.